

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Frequency distribution of six cytokine gene polymorphisms in North- and South-Italy

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1661329> since 2018-04-19T11:35:08Z

Published version:

DOI:10.1111/iji.12324

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



Frequency distribution of six cytokine gene polymorphisms in North- and South-Italy -Italy

Journal:	<i>International Journal of Immunogenetics</i>
Manuscript ID	IJIG-Feb-17-0022.R1
Manuscript Type:	Short Communication
Date Submitted by the Author:	10-Mar-2017
Complete List of Authors:	Santovito, Alfredo; Universita degli Studi di Torino, Department of Life Sciences and Systems Biology; Gendusa, Claudio; Universita degli Studi di Torino, Department of Life Sciences and Systems Biology Matini, Alessandro; Universita degli Studi di Torino, Department of Life Sciences and Systems Biology Ferraro, Francesca; Universita degli Studi di Torino, Department of Life Sciences and Systems Biology Musso, Irene; Universita degli Studi di Torino, Department of Life Sciences and Systems Biology Costanzo, Maria; Universita degli Studi di Torino, Department of Life Sciences and Systems Biology Delcos, Amandine; Universita degli Studi di Torino, Department of Life Sciences and Systems Biology Cervella, Piero; Universita degli Studi di Torino, Department of Life Sciences and Systems Biology
Keywords:	Cytokine, population studies < polymorphism < Cytokine, allele frequencies < molecular < Genetics, gene - polymorphism < DNA < Molecular Biology

Short Communications

Population Study

Frequency distribution of six cytokine gene polymorphisms in North- and South-Italy

Running Head: Cytokine gene polymorphisms in an Italian sample

Alfredo SANTOVITO*, Claudio GENDUSA, Alessandro MATINI, Francesca FERRARO, Irene MUSSO, Maria COSTANZO, Amandine DELCLOS, and Piero CERVELLA

University of Turin (Italy), Department of Life Sciences and Systems Biology

*Corresponding Author:

Alfredo SANTOVITO

Department of Life Sciences and Systems Biology

Via Accademia Albertina n. 13 - 10123 – Torino (Italy)

Tel.: +39-0116704554; Fax: +39-0116704508;

e-mail: alfredo.santovito@unito.it

Keywords: Cytokines; Population Study; Allele Frequencies; Gene polymorphism

Summary

Allelic and genotype frequencies of four cytokine genes were obtained from 738 subjects from North- and South-Italy. Populations were in Hardy-Weinberg equilibrium for all genes but significantly differed in the frequency of all SNPs and three haplotypes. In the MDS graph, they were plotted in separate positions close to Europeans and an Ivorian population, respectively.

Introduction

Cytokines are small regulatory proteins mainly secreted by active immune cells in response to different stimuli such as infection and tissue damage. These molecules play a crucial role in regulating all aspects of immune and inflammatory responses and represent key components in the pathogenesis of many diseases like cancer, metabolic disorders, infectious and autoimmune diseases (for reviews see: Bidwell *et al.*, 1999, 2001; Haukim *et al.*, 2002; Hollegaard & Bidwell, 2006).

Inter-individual differences in the related cytokines serum levels were observed (Hoffmann *et al.*, 2002). These differences in cytokine production and, consequently, in the individual response to various antigens were attributed to a different factors, including gene polymorphisms. Indeed, certain cytokine polymorphisms, mostly single nucleotide polymorphisms (SNPs) located within in the promoter or coding regions, have been shown to affect the overall expression and secretion of the gene products (Hoffmann *et al.*, 2001).

At individual level, this different cytokines expression could explain the different susceptibility to various diseases including autoimmune, infectious or cancer diseases observed among individual belonging to the same population. At population level, it was found that the distribution of cytokine polymorphisms significantly varies among different ethnic groups, mainly as a result of founder effects and geographically localized selective forces (Santovito *et al.*, 2012; Hollegaard & Bidwell, 2006; Hoffmann *et al.*, 2002). For this reason, the association of cytokine gene polymorphisms with a particular disease cannot be extrapolated from a specific population to other populations with

different genetic background. On the other hand, data concerning distribution of cytokine polymorphisms in healthy populations are important in order to investigate the possible associations of these polymorphisms with a particular disease, especially in case-control studies.

Finally, the analysis of cytokine gene polymorphisms in different populations has been used in anthropological studies in order to establish possible gradient in the distribution of the genetic variation among human populations.

Italy has received the passage of multiple human groups in prehistoric and historic times: Phoenician, Greek, Carthaginian, Roman, Arabic, Norman and Barbaric populations contributed to the present genetic composition of Italy (Rickards et al., 1998). Genome wide association studies evidenced a genetic structure of the Italian population strongly influenced by the geographical distance, with certain degree of genetic substructure between Southern Italians, Northern Italy and other European populations (Nelis et al. 2009; Di Gaetano et al., 2012). In particular, studies based on mtDNA and Y-chromosome variability (Barbujani et al., 1995; Capelli et al., 2007) identified a North-South gradient of the genetic variation within the peninsula. Differential Neolithic/Mesolithic contributes in the two regions as well as local drift and founder effects were invoked to explain the observed genetic variation distribution (Cappelli et al., 2007; Di Giacomo et al., 2003).

In this scenario, the aim of the present study was to evaluate the alleles, genotypes and haplotypes frequencies of selected cytokine gene polymorphisms in South and North Italian populations, and to compare these allele frequencies with those already published for other populations worldwide distributed.

Materials and Methods

Subjects

The study was conducted on a cohort of 738 unrelated healthy Italian subjects of Caucasian origin collected specifically for this epidemiological study and not for diseases studies: 635 from North

Italy (359 females and 276 males, mean age 51.04 ± 0.87 , 341 from Piedmont, 198 from Lombardy and 96 from Aosta regions) and 103 from South Italy (Sicily region, 53 females and 50 males, mean age 49.63 ± 1.65 , 62 from Trapani and 41 from Pantelleria Island). All the subjects were randomly chosen healthy volunteers, received detailed information about the study, and gave their informed consent prior the analyses.

The study was approved by the University of Turin ethics committee and was performed in agreement with the ethical standards laid down in the 1964 Declaration of Helsinki.

DNA extraction and Genotyping

Peripheral blood samples (5-10 ml obtained by venipuncture) were collected in heparinized vacutainers and stored at -20°C . DNA extraction was conducted using a Chelex solution, according to the following protocol: 10 μL of peripheral blood was diluted in 1 mL of sterile distilled water for 15 min at room temperature. After centrifugation at 14,000 rpm for 1 min, the pellet was re-suspended in 200 μL of 5% Chelex solution in Tris-EDTA at pH 8, heated to 56°C for 15 min and, after vortex for 10 sec, at 100°C in boiler water for 8 min. For PCR reactions we used 19 of the 200 μL of this solution, containing about 10 ng of DNA as indicated by the spectrophotometric analysis. PCR-based genotyping was performed for the genes encoding *TNF α* (G/A -308), *IL10* (G/A -1082, C/T -819), *TGF β* (C/T codon 10, G/C codon 25), *IL6* (G/C -174). The sequence polymorphisms were determined by SSP-PCR methodology, using primers described in Perrey et al., (1999). PCR reactions were performed in a 25 μL volume, with a final concentration of 1X Reaction Buffer, 1.5 mM of MgCl_2 , 5% of DMSO, 250 μM of dNTPs, 0.5 μM of each primer, and 1 U/sample of Taq DNA polymerase (Fischer, U.S.). Cycles were set as follows: 35 cycles, 1 min at 95°C , 1 min at 60°C , 1 min at 72°C , and a final extension step 10 min at 72°C . Amplification products were detected by ethidium bromide staining after 2.5% agarose gel electrophoresis. To verify the genotyping results, about 10% of the total sample ($n = 80$) were also analysed by another investigator. The two analyses showed identical results.

Statistical Analysis

Allele, genotype and haplotype frequencies of the six SNPs were calculated by direct counting, dividing by the number of subjects (to produce genotype frequency) or chromosomes (to produce allele and haplotype frequency). The Pearson chi-square test (χ^2 -test), with a 95% confidence interval, as implemented in SPSS statistical package program (v23.0, IBM, Chicago, IL), was used to analyse possible statistical differences in allele and haplotype frequencies between studied populations, as well as to test the Hardy-Weinberg equilibrium (HWE). Allele frequencies found in our populations were compared with published data for other worldwide distributed populations. Genetic relationships among these populations were analysed via non-metric multidimensional scaling (MDS) analysis of allele frequencies, as implemented in SPSS, and by plotting the positions of the populations using the two principal dimensions. We computed the Euclidean distance for each pair of populations using cytokine allele frequencies. This dissimilarity matrix was used to generate a MDS plot of population variation in two dimensions.

Results and Discussion

Allele and genotype frequencies of North and South Italian populations were reported in Table 1. In both studied populations the genotype frequency distribution of all polymorphic cytokine genes did not show a significant deviation from HWE, indicating a random distribution and the absence of evolutionary forces acting in shaping the frequencies of these gene polymorphisms. However, when we compared the two populations, we found significant differences for all SNPs of the cytokine polymorphisms tested. Therefore, we considered these two Italian populations as two separate groups when compared with other populations.

We further calculated the haplotype frequencies of *IL10* (-1082G/A, -819C/T) and *TGFβ1* (cod.10 C/G, cod. 25 C/T) (Table 2). For *IL10* (-1082G/A, -819C/T), a total of 4 haplotypes (for both North- and South-Italy, samples) was found, with the GC haplotype showing the higher frequency in both

populations (0.796 and 0.602 for North- and South-Italy, respectively), followed by the GT haplotype (0.090 and 0.199 for North- South-Italy, respectively). *IL10* (-1082G/A, -819C/T) GG/CC was the most common genotype found, with a frequency of about 0.613 and 0.359 for North- and South-Italian populations, respectively, while the least common genotype was GA/TT for North-Italy (0.002) and AA/CC for South-Italy where it resulted absent.

The most common *TGFβ1* (cod.10 C/G, cod. 25 C/T) haplotype was CG for both populations (with frequencies of 0.765 and 0.840 for North- and South-Italy, respectively), followed by TG haplotype found at 0.162 and 0.082 in North- and South-Italian populations, respectively. The most common *TGFβ1* (cod.10 C/G, cod. 25 C/T) genotype was CC/GG, with a frequency of 0.586 in North-Italy and 0.728 in South-Italy, while the least common was CC/CC (0.003) for North-Italy and TT/GG (0.010) for South-Italy. Significant differences were found between North- and South-Italy in the frequency of *IL10* (-1082G/A, -819C/T) GC ($P = 0.021$), GT ($P < 0.001$) and AT ($P = 0.002$) haplotypes as well as in the frequency of *TGFβ1* (cod.10 C/G, cod. 25 C/T) TG haplotype ($P = 0.008$).

In order to further assess population relationships, a MDS analysis based on the cytokine allele frequencies was carried out comparing our studied populations with some worldwide distributed populations data set typed for the same loci (Nancy et al., 2004; Louie et al., 2005; Mihailova et al., 2005; Trajkov et al., 2005; Kubistova et al., 2006; Skorpil et al., 2007; Javor et al., 2007; Sodsai et al., 2011; Visentainer et al., 2008; Norhalifah et al., 2015; Santovito et al., 2012; Kaur et al., 2007; Costeas et al., 2003) (Fig. 1). MDS can be considered to be an alternative to factor analysis that allows to analyse any kind of similarity or dissimilarity matrix, in addition to correlation matrices. In this case, the two dimensions account for 99% of the observed variance and were used to plot the positions of the populations.

In the MDS plot, the studied Italian populations were separated from the major group: North-Italy was plotted in the same quadrant of other European populations, such as Slovak, Czech, Germany, Netherland and Macedonian populations, although separated from them. South-Italy, *vice versa*,

was well separated from all the other populations and plotted in the same quadrant of the Ivory Coast population.

In general, when compared to other European populations, the Northern Italian population was genetically close to North-European populations, such as the French population, whereas the Southern Italians had some similarities with other Mediterranean populations, as well as with those from Middle East (Di Gaetano et al., 2012).

Our results seem to be concordant with data obtained by other authors with more polymorphic markers, showing for Italy a differential genetic diversity pattern between North- and South-Italy (Boattini et al., 2013). This picture probably reflects the different genetic history of Sicily with respect to Northern Italy. Indeed, because to its central geographic location in the Mediterranean domain, Sicily has long been the meeting place of ancient civilizations and cultures and hosted various human groups in both prehistoric and historic times (Sarno et al., 2014). As consequence, various populations have contributed to the genetic structure of this island. Since the arrival of the first human groups (Sicani, Siculi, Elymi), the island has been subjected to numerous migratory flows: Greeks, Phoenicians, Etruscans, Romans, Vandals, Goths, Byzantines, Arabs, Aragoneses and Normans contribute to the strong heterogeneity observed in the genetic structure of Sicilian populations (Rickards et al. 1998; Cerutti et al., 2004).

For other populations, such as Asian and African populations, the observed differences could be explained by founder effects and/or local selective pressures imposed by host-pathogen interactions on specific geographic populations (Hollegaard & Bidwell, 2006; Santovito et al., 2012). Finally, for the European populations differences could be probably explained with the different sample size and possible stochastic factors, although also for these populations micro-evolutionary forces cannot be excluded.

Conclusion

In conclusion, in this study we found that the frequencies of analyzed cytokine alleles showed a significant different distribution between North- and South-Italy, as also evidenced in the MDS plot. Because of the ethnical and local differences in the distribution of cytokine gene polymorphisms, the population data from healthy individuals are of relevant interest for the evaluation of the role of these polymorphisms in the differential response to various immunological diseases and in the occurrence of those diseases influenced by variations of cytokines production.

Ethics

Samples included in present study were collected after informed consent.

Funding

The study was supported by grants (named “ex 60%”) from the Italian Ministry of University and Scientific Research.

Disclosures

None of the authors have a conflict of interest to declare in relation to this work.

References

- Barbujani, G., Bertorelle, G., Capitani, G. & Scozzari, R. (1995) Geographical structuring in the mtDNA of Italians. *Proc Natl Acad Sci U S A*, 92, 9171–9175.
- Bidwell, J., Keen, L., Gallagher, G., Kimberly, R., Huizinga, T., & McDermott, M.F., et al. (1999). Cytokine gene polymorphism in human disease: on-line databases. *Genes and Immunity*, 1, 3-19. doi: 10.1038/sj.gene.6363645

Bidwell, J., Keen, L., Gallagher, G., Kimberly, R., Huizinga, T., McDermott, M.F., et al. (2001). Cytokine gene polymorphism in human disease: on-line databases, supplement 1. *Genes and Immunity*, 2, 61-70. doi: 10.1038/sj.gene.6363733

Boattini, A., Martinez-Cruz, B., Sarno, S., Harmat, C., Useli, A., Sanz, P., et al. (2013) Uniparental Markers in Italy Reveal a Sex-Biased Genetic Structure and Different Historical Strata. *PLoS One*, 8, e65441. doi: 10.1371/journal.pone.0065441

Capelli, C., Brisighelli, F., Scarnicci, F., Arredi, B., Caglia, A., Vetrugno, G. et al. (2007) Y chromosome genetic variation in the Italian peninsula is clinal and supports an admixture model for the Mesolithic-Neolithic encounter. *Molecular Phylogenetics and Evolution*, 44, 228–239. doi: 10.1016/j.ympev.2006.11.030

Cerutti, N., Dugoujon, J.M., Guitard, E. & Rabino Massa, E. (2004). Gm and Km immunoglobulin allotypes in Sicily. *Immunogenetics*, 55, 674-681. doi: 10.1007/s00251-003-0628-z

Costeas, P.A., Koumas, L., Koumouli, A., Kyriakou-Giantsiou, A. & Papaloizou Haukim, A. (2003). Cytokine polymorphism frequencies in the Greek Cypriot population. *European Journal of Immunogenetics*, 30, 341-343. doi: 10.1046/j.1365-2370.2003.00413.x

Di Gaetano, C., Voglino, F., Guarrera, S., Fiorito, G., Rosa, F., Di Blasio, A.M. et al. (2012) An Overview of the Genetic Structure within the Italian Population from Genome-Wide Data. *PLoS One*, 7, e43759. doi: 10.1371/journal.pone.0043759

Di Giacomo, F., Luca, F., Anagnou, N., Ciavarella, G., Corbo, R.M., Cresta, M. et al. (2003) Clinal patterns of human Y chromosomal diversity in continental Italy and Greece are dominated by drift and founder effects. *Molecular Phylogenetics and Evolution* 28, 387–395. doi: 10.1016/S1055-7903(03)00016-2

- Haukim, N., Bidwell, J.L., Smith A.J., Keen, L.J., Gallagher, G., Kimberly, R., et al. (2002). Cytokine gene polymorphism in human disease: on-line databases, supplement 2. *Genes and Immunity*, 3(6), 313-330. doi:10.1038/sj.gene.6363881
- Hoffmann, S.C., Stanley, E.M., Darrin, C.E., Craighead, N., DiMercurio, B.S., Koziol, D.E. et al. (2001). Association of cytokine polymorphic inheritance and in vitro cytokine production in anti-CD3/CD28-stimulated peripheral blood lymphocytes. *Transplantation*, 72, 1444-1450.
- Hoffmann, S.C., Stanley, E.M., Cox, E.D., DiMercurio, B.S., Koziol, D.E., Harlan, D.M. et. al. (2002). Ethnicity greatly influences cytokine gene polymorphism distribution. *American Journal of Transplantation*, 2, 560-567. doi: 10.1034/j.1600-6143.2002.20611.x
- Hollegaard, M.V. & Bidwell, J.L. (2006). Cytokine gene polymorphism in human disease: on-line databases, Supplement 3. *Genes & Immunity*, 7(4), 269-276. doi:10.1038/sj.gene.6364301.
- Javor, J., Bucova, M., Ferencik, S., Grosse-Wilde H. & Buc, M. (2007). Single nucleotide polymorphisms of cytokine genes in the healthy Slovak population. *International Journal of Immunogenetics*, 34, 273-280. doi: 10.1111/j.1744-313X.2007.00693x
- Kaur, G., Rapthap, C.C., Kumar, N., Kumar, S., Neolia S. & Mehra, N.K. (2007). Frequency distribution of cytokine gene polymorphisms in the healthy North Indian population. *Tissue Antigens*, 69, 113-120. doi: 10.1111/j.1399-0039.2006.00740.x
- Kubistova, Z., Mrazek, F., Tudos, Z., Kriegova, E., Ambrozova, Z., Mytilineos, J. & Petrek, M. (2006). Distribution of 22 cytokine gene polymorphisms in the healthy Czech population. *International Journal of Immunogenetics*, 33, 261-267. doi: 10.1111/j.1744-313X.2006.00609.x
- Louie, L.G., Silver, E.W., Direskeneli, G.S., Kearney, F.C., Spiroski, M.Z., Peste-Tsilimidou, C. et al. (2005) Worldwide variation in cytokine genes. In: HLA 2002 - Immunobiology of the Human MHC (ed. by J. A. Hansen & B. Dupont). IHWG Press, Seattle, 2005.
- Mihailova, S., Ivanova, M., Mihaylova, A., Quin, L., Mikova, O. & Naumova, E. (2005) Pro- and anti-inflammatory cytokine gene polymorphism profiles in Bulgarian multiple sclerosis patients. *Journal of Neuroimmunology*, 168, 138. doi:10.1016/j.jneuroim.2005.06.020

Nancy, L., Delaney, N.L., Esquenazi, V., Lucas, D.P., Zachary, A.A. & Leffell, M.S. (2004). TNF- α , TGF- β , IL-10, IL-6, and INF- γ Alleles Among African Americans and Cuban Americans. Report of the ASHI. Minority Workshops: Part IV. Human Immunology, 65, 1413-1419. doi: 10.1016/j.humimm.2004.07.240

Nelis, M., Esko, T., Magi, R., Zimprich, F., Zimprich, A., Tonchevaet, D. et. al. (2009) Genetic structure of Europeans: a view from the North-East. *PLoS One*, 4: e5472. doi: 10.1371/journal.pone.0005472

Norhalifah, H. K., Zafarina, Z., Sundararajulu, P., Norazmi, M.N. & Edinur, H.A. (2015) Distribution of cytokine gene polymorphisms in five Malay subethnic groups in Peninsular Malaysia. *International Journal of Immunogenetics*, 42, 200-203. doi: 10.1111/iji.12189

Perrey, C., Turner, S.J., Pravica, V., Howell, W.M. & Hutchinson, I.V. (1999). ARMS-PCR methodologies to determine IL-10, TNF- α , TNF- β and TGF- β 1 gene polymorphisms. *Transplant Immunology*, 7, 127-128. doi: 10.1016/S0966-3274(99)80030-6

Rickards, O., Martinez-Labarga, C., Scano, G., De Stefano, G.F., Biondi, G.F., Pacaci, M. & Walter H (1998) Genetic history of the population of Sicily. *Human Biology*, 70, 699–714.

Santovito, A., Cervella, P, Schleicherova, D, & Delpero, M. (2012). Genotyping for cytokine polymorphisms in a Northern Ivory Coast population reveals a high frequency of the heterozygote genotypes for the TNF-a-308G/A SNP. *International Journal of Immunogenetics*, 39, 291-295. doi: 10.1111/j.1744-313X.2012.01086.x

Sarno, S., Boattini, A., Carta, M., Gianmarco, F., Alù, M., Yang Yao, et al. (2014) An Ancient Mediterranean Melting Pot: Investigating the Uniparental Genetic Structure and Population History of Sicily and Southern Italy. *PLoS One*, 9, e96074. doi:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[10.1371/journal.pone.0096074](https://doi.org/10.1371/journal.pone.0096074)

Skorpil, N., Kolesár, L., Striz, I., Lardy, N.M. & Slavcev, A. (2007) Cytokine gene polymorphisms in the Dutch population. *International Journal of Immunogenetics*, 34, 87-90. doi: 10.1111/j.1744-313X.2007.00663.x

Sodsai, P., Nakkuntod, J., Kupatawintu, P. & Hirankarn, N. (2011) Distribution of cytokine gene polymorphisms in Thai Population. *Tissue Antigens*, 77, 593-597. doi: 10.1111/j.1399-0039.2011.01647.x


Trajkov, D., Arsov, T., Petlichovski, A., Strezova, A., Efinska-Mladenovska, O. & Spiroski, M. (2005) Cytokine gene polymorphisms in population of ethnic Macedonians. *Croatian Medical Journal*, 46, 685.

Visentainer, J.E.L., Sell, A.M., de Silva, G.C., Cavicchioli, A.D.G., Franceschi, D.S.A., Lieber, S.R., & de Souza, C.A. (2008). TNF, IFNG, IL6, IL10 and TGFB1 gene polymorphisms in South and Southeast Brazil. *International Journal of Immunogenetics*, 35, 287-293. doi: 10.1111/j.1744-313X.2008.00778.x

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1 - Nonmetric MDS applied in R matrix based on six cytokine gene polymorphisms analyzing the genetic relationships among some populations worldwide distributed. The two axes explain about 99% of the variation in allele frequencies.

For Peer Review



223x184mm (72 x 72 DPI)

Table 1 – Allele and Genotype Frequencies of six cytokine gene polymorphisms In North- (n = 635) and South-Italy (n = 103)

Cytokine polymorphisms	Allele	N	Frequency	Genotype	N	Frequency	HWE P-value
TNFA-308 North-Italy	A	155	0.122	AA	6	0.009	0.440
	G	1115	0.878	AG	143	0.225	
				GG	486	0.765	
	A	59	0.286	AA	6	0.058	0.498
	G	147	0.714	AG	47	0.456	
				GG	50	0.485	
IL6 -174 North-Italy	C	114	0.090	CC	5	0.008	0.998
	G	1156	0.910	CG	104	0.164	
				GG	526	0.828	
	C	48	0.233	CC	9	0.621	0.171
	G	158	0.767	CG	30	0.291	
				GG	64	0.087	
IL10 -1082 North-Italy	A	144	0.133	AA	6	0.009	0.694
	G	1126	0.887	AG	132	0.208	
				GG	497	0.783	
	A	41	0.199	AA	4	0.039	0.999
	G	165	0.801	AG	33	0.320	
				GG	66	0.641	
IL10 -819 North-Italy	T	155	0.122	TT	8	0.013	0.864
	C	1115	0.878	CT	139	0.219	
				CC	488	0.769	
	T	58	0.282	TT	12	0.117	0.175
	C	148	0.718	CT	34	0.330	
				CC	57	0.533	
TGFB-codon 10 North-Italy	T	233	0.183	CC	28	0.044	0.214
	C	1037	0.817	CT	177	0.279	
				TT	430	0.677	
	T	24	0.117	CC	3	0.029	0.309
	C	182	0.883	CT	18	0.175	
				TT	82	0.796	
TGFB-codon 25 North-Italy	C	93	0.073	CC	2	0.003	0.714
	G	1177	0.927	CG	89	0.140	
				GG	544	0.857	
	C	16	0.078	CC	2	0.019	0.166
	G	190	0.922	CG	12	0.117	
				GG	89	0.864	

HWE = Hardy-Weinberg Equilibrium. Significant differences for all SNPs of the cytokine polymorphisms tested were found between North- and South-Italy.

Table 2 – *IL10* (-1082G/A, -819C/T) and *TGFβ1* (cod.10 C/G, cod. 25 C/T) genotype and haplotype frequency in the studied North- (n = 635) and South- (n = 103) Italian populations.

Cytokine	North-Italy N (frequency)	South-Italy N (frequency)	$P \chi^2$ - test
<i>IL10</i> (-1082G/A, -819C/T)			
Genotype			
GG/CC	389 (0.613)	37 (0.359)	0.009
GG/CT	101 (0.159)	21 (0.204)	0.333
GG/TT	7 (0.011)	8 (0.078)	0.001
GA/CC	95 (0.150)	20 (0.194)	0.320
GA/CT	36 (0.057)	9 (0.087)	0.272
GA/TT	1 (0.002)	4 (0.039)	0.002
AA/CC	4 (0.006)	0 (0.000)	1.000
AA/CT	2 (0.003)	4 (0.039)	0.005
<i>IL10</i> (-1082G/A, -819C/T)			
Haplotypes			
GC	1010 (0.796)	24 (0.602)	0.021
GT	114 (0.090)	41 (0.199)	<0.001
AC	105 (0.083)	24 (0.116)	0.151
AT	39 (0.031)	17 (0.083)	0.002
<i>TGFβ1</i> (cod.10 C/G, cod. 25 C/T)			
Genotype			
CC/GG	372 (0.586)	75 (0.728)	0.208
CC/GC	50 (0.079)	5 (0.049)	0.415
CC/CC	2 (0.003)	2 (0.019)	0.098
CT/GG	143 (0.225)	13 (0.126)	0.066
CT/GC	34 (0.054)	5 (0.049)	1.000
TT/GG	29 (0.046)	1 (0.010)	0.107
TT/GC	5 (0.008)	2 (0.019)	0.258
<i>TGFβ1</i> (cod.10 C/G, cod. 25 C/T)			
Haplotype			
CG	971 (0.765)	173 (0.840)	0.402
TG	206 (0.162)	17 (0.082)	0.008
CC	54 (0.042)	9 (0.044)	0.854
TC	39 (0.031)	7 (0.034)	0.828

$P \chi^2$ = Probability from Chi-square test between North and South Italy

Significant results are highlighted in gray



COPYRIGHT TRANSFER AGREEMENT

Date: 02-02-2017 Contributor name: Alfredo SANTOVITO
Contributor address: Department of Life Sciences and Systems Biology, Via Accademia Albertina 13, 10123 Turin (Italy)
Manuscript number (if known):
Re: Manuscript entitled Frequency distribution of six cytokine gene polymorphisms in a sample of healthy subjects
from North- and South-Italy (the "Contribution")
for publication in International Journal of Immunogenetics (the "Journal")
published by Wiley-Blackwell ("Wiley-Blackwell").

Dear Contributor(s):
Thank you for submitting your Contribution for publication. In order to expedite the editing and publishing process and enable Wiley-Blackwell to disseminate your Contribution to the fullest extent, we need to have this Copyright Transfer Agreement signed and returned as directed in the Journal's instructions for authors as soon as possible. If the Contribution is not accepted for publication, or if the Contribution is subsequently rejected, this Agreement shall be null and void. **Publication cannot proceed without a signed copy of this Agreement.**

A. COPYRIGHT

1. The Contributor assigns to Wiley-Blackwell, during the full term of copyright and any extensions or renewals, all copyright in and to the Contribution, and all rights therein, including but not limited to the right to publish, republish, transmit, sell, distribute and otherwise use the Contribution in whole or in part in electronic and print editions of the Journal and in derivative works throughout the world, in all languages and in all media of expression now known or later developed, and to license or permit others to do so.
2. Reproduction, posting, transmission or other distribution or use of the final Contribution in whole or in part in any medium by the Contributor as permitted by this Agreement requires a citation to the Journal and an appropriate credit to Wiley-Blackwell as Publisher, and/or the Society if applicable, suitable in form and content as follows: (Title of Article, Author, Journal Title and Volume/Issue, Copyright © [year], copyright owner as specified in the Journal). Links to the final article on Wiley-Blackwell's website are encouraged where appropriate.

B. RETAINED RIGHTS

Notwithstanding the above, the Contributor or, if applicable, the Contributor's Employer, retains all proprietary rights other than copyright, such as patent rights, in any process, procedure or article of manufacture described in the Contribution.

C. PERMITTED USES BY CONTRIBUTOR

1. **Submitted Version.** Wiley-Blackwell licenses back the following rights to the Contributor in the version of the Contribution as originally submitted for publication:
a. After publication of the final article, the right to self-archive on the Contributor's personal website or in the Contributor's institution's/employer's institutional repository or archive. This right extends to both intranets and the Internet. The Contributor may not update the submission version or replace it with the published Contribution. The version posted must contain a legend as follows: This is the pre-peer reviewed version of the following article: FULL CITE, which has been published in final form at [Link to final article].
b. The right to transmit, print and share copies with colleagues.
2. **Accepted Version.** Re-use of the accepted and peer-reviewed (but not final) version of the Contribution shall be by separate agreement with Wiley-Blackwell. Wiley-Blackwell has agreements with certain funding agencies governing reuse of this version. The details of those relationships, and other offerings allowing open web use, are set forth at the following website: <http://www.wiley.com/go/funderstatement>. NIH grantees should check the box at the bottom of this document.

3. **Final Published Version.** Wiley-Blackwell hereby licenses back to the Contributor the following rights with respect to the final published version of the Contribution:

- a. Copies for colleagues. The personal right of the Contributor only to send or transmit individual copies of the final published version in any format to colleagues upon their specific request provided no fee is charged, and further-provided that there is no systematic distribution of the Contribution, e.g. posting on a listserve, website or automated delivery.
- b. Re-use in other publications. The right to re-use the final Contribution or parts thereof for any publication authored or edited by the Contributor (excluding journal articles) where such re-used material constitutes less than half of the total material in such publication. In such case, any modifications should be accurately noted.
- c. Teaching duties. The right to include the Contribution in teaching or training duties at the Contributor's institution/place of employment including in course packs, e-reserves, presentation at professional conferences, in-house training, or distance learning. The Contribution may not be used in seminars outside of normal teaching obligations (e.g. commercial seminars). Electronic posting of the final published version in connection with teaching/training at the Contributor's institution/place of employment is permitted subject to the implementation of reasonable access control mechanisms, such as user name and password. Posting the final published version on the open Internet is not permitted.
- d. Oral presentations. The right to make oral presentations based on the Contribution.

4. **Article Abstracts, Figures, Tables, Data Sets, Artwork and Selected Text (up to 250 words).**

- a. Contributors may re-use unmodified abstracts for any non-commercial purpose. For on-line uses of the abstracts, Wiley-Blackwell encourages but does not require linking back to the final published versions.
- b. Contributors may re-use figures, tables, data sets, artwork, and selected text up to 250 words from their Contributions, provided the following conditions are met:
 - (i) Full and accurate credit must be given to the Contribution.
 - (ii) Modifications to the figures, tables and data must be noted. Otherwise, no changes may be made.
 - (iii) The reuse may not be made for direct commercial purposes, or for financial consideration to the Contributor.
 - (iv) Nothing herein shall permit dual publication in violation of journal ethical practices.

D. CONTRIBUTIONS OWNED BY EMPLOYER

1. If the Contribution was written by the Contributor in the course of the Contributor's employment (as a "work-made-for-hire" in the course of employment), the Contribution is owned by the company/employer which must sign this Agreement (in addition to the Contributor's signature) in the space provided below. In such case, the company/employer hereby assigns to Wiley-Blackwell, during the full term of copyright, all copyright in and to the Contribution for the full term of copyright throughout the world as specified in paragraph A above.

2. In addition to the rights specified as retained in paragraph B above and the rights granted back to the Contributor pursuant to paragraph C above, Wiley-Blackwell hereby grants back, without charge, to such company/employer, its subsidiaries and divisions, the right to make copies of and distribute the final published Contribution internally in print format or electronically on the Company's internal network. Copies so used may not be resold or distributed externally. However the company/employer may include information and text from the Contribution as part of an information package included with software or other products offered for sale or license or included in patent applications. Posting of the final published Contribution by the institution on a public access website may only be done with Wiley-Blackwell's written permission, and payment of any applicable fee(s). Also, upon payment of Wiley-Blackwell's reprint fee, the institution may distribute print copies of the published Contribution externally.

E. GOVERNMENT CONTRACTS

In the case of a Contribution prepared under U.S. Government contract or grant, the U.S. Government may reproduce, without charge, all or portions of the Contribution and may authorize others to do so, for official U.S. Govern-

ment purposes only, if the U.S. Government contract or grant so requires. (U.S. Government, U.K. Government, and other government employees: see notes at end)

F. COPYRIGHT NOTICE

The Contributor and the company/employer agree that any and all copies of the final published version of the Contribution or any part thereof distributed or posted by them in print or electronic format as permitted herein will include the notice of copyright as stipulated in the Journal and a full citation to the Journal as published by Wiley-Blackwell.

G. CONTRIBUTOR'S REPRESENTATIONS

The Contributor represents that the Contribution is the Contributor's original work, all individuals identified as Contributors actually contributed to the Contribution, and all individuals who contributed are included. If the Contribution was prepared jointly, the Contributor agrees to inform the co-Contributors of the terms of this Agreement and to obtain their signature to this Agreement or their written permission to sign on their behalf. The Contribution is submitted only to this Journal and has not been published before. (If excerpts from copyrighted works owned by third parties are included, the Contributor will obtain written permission from the copyright owners for all uses as set forth in Wiley-Blackwell's permissions form or in the Journal's Instructions for Contributors, and show credit to the sources in the Contribution.) The Contributor also warrants that the Contribution contains no libelous or unlawful statements, does not infringe upon the rights (including without limitation the copyright, patent or trademark rights) or the privacy of others, or contain material or instructions that might cause harm or injury.

CHECK ONE BOX:

☒ Contributor-owned work

ATTACH ADDITIONAL SIGNATURE
PAGES AS NECESSARY

Contributor's signature

Alfredo Santovito

Date 02-02-2017

Type or print name and title Alfredo SANTOVITO, PhD, Dr

Co-contributor's signature

Date

Type or print name and title

☐ Company/Institution-owned work
(made-for-hire in the
course of employment)

Company or Institution (Employer-for-Hire)

Date

Authorized signature of Employer

Date

☐ U.S. Government work

Note to U.S. Government Employees

A contribution prepared by a U.S. federal government employee as part of the employee's official duties, or which is an official U.S. Government publication, is called a "U.S. Government work," and is in the public domain in the United States. In such case, the employee may cross out Paragraph A.1 but must sign (in the Contributor's signature line) and return this Agreement. If the Contribution was not prepared as part of the employee's duties or is not an official U.S. Government publication, it is not a U.S. Government work.

☐ U.K. Government work
(Crown Copyright)

Note to U.K. Government Employees

The rights in a Contribution prepared by an employee of a U.K. government department, agency or other Crown body as part of his/her official duties, or which is an official government publication, belong to the Crown. U.K. government authors should submit a signed declaration form together with this Agreement. The form can be obtained via <http://www.opsi.gov.uk/advice/crown-copyright/copyright-guidance/publication-of-articles-written-by-ministers-and-civil-servants.htm>

☐ Other Government work

Note to Non-U.S., Non-U.K. Government Employees

If your status as a government employee legally prevents you from signing this Agreement, please contact the editorial office.

☐ NIH Grantees

Note to NIH Grantees

Pursuant to NIH mandate, Wiley-Blackwell will post the accepted version of Contributions authored by NIH grant-holders to PubMed Central upon acceptance. This accepted version will be made publicly available 12 months after publication. For further information, see www.wiley.com/go/nihamandate.